arXiv:1403.7573v1 [math.DG] 29 Mar 2014

ON THE BIHARMONIC CURVES IN THE SPECIAL LINEAR GROUP $\mathrm{SL}(2,\mathbb{R})$

I. I. ONNIS AND A. PASSOS PASSAMANI

Abstract. We characterize the biharmonic curves in the special linear group $\mathrm{SL}(2,\mathbb{R})$. In particular, we show that all proper biharmonic curves in $\mathrm{SL}(2,\mathbb{R})$ are helices and we give their explicit parametrizations as curves in the pseudo-Euclidean space $\mathbb{R}^4_2$.

1. Introduction

Let $\phi : (M^m, g) \to (N^n, h)$ be a smooth map between two Riemannian manifolds. The tension field of ϕ is, by definition, $\tau(\phi) = \operatorname{trace} \nabla^N d(\phi)$. According J. Eells and J.H. Sampson, see [8], ϕ is biharmonic if it is a critical point of the bienergy functional

$$E_2(\phi) = \frac{1}{2}\int_M |\tau(\phi)|^2 v_g. \tag{1}$$

The first variation formula for E_2 was compute by G.Y. Jiang in [9] and [10] as

$$\tau_2(\phi) := -\Delta^\phi \tau(\phi) - \operatorname{trace} \mathrm{R}^N(d\phi, \tau(\phi))d\phi = 0, \tag{2}$$

where Δ^ϕ denotes the *rough Laplacian* acting on $C(\phi^{-1}TN)$, defined by

$$\Delta^\phi = -\operatorname{trace}(\nabla^\phi)^2 = -\sum_{i=1}^{m}\{\nabla^\phi_{E_i}\nabla^\phi_{E_i} - \nabla^\phi_{\nabla^M_{E_i}E_i}\}. \tag{3}$$

with respect to a local orthonormal frame field $\{E_i\}_{i=1}^m$ on M.
The field $\tau_2(\phi)$ is named *bitension field* of ϕ.

As a geodesic curve ($\tau(\phi) = 0$) is a biharmonic one, we are interested in biharmonic curves that are not geodesics i.e. *proper biharmonic curves*.

The study of the proper biharmonic curves on a curved surface starts with [5] where there are described these curves in a surface, proving that biharmonic curves on a surface of non-positive Gaussian curvature are geodesics.
For 3-dimensional Riemannian manifolds with constant sectional curvature, the case of null and negative curvature are considered in [7] and [3] and it is showed that the only biharmonic curves are the geodesic ones. Moreover, in [2], it is considered the case of positive curvature showing that biharmonic curves have constant geodesic curvature and geodesic torsion.

Date: March 2014.
1991 *Mathematics Subject Classification.* 53C30, 58E20.
Key words and phrases. Special linear group, biharmonic curves, helix curves, homogeneous spaces.
The second author was supported by Capes–Brazil.

Besides the spaces forms, the most relevant 3-dimensional homogeneous Riemannian spaces are those with 4-dimensional isometry group: the Berger spheres, the Heisenberg group, the special linear group $\mathrm{SL}(2,\mathbb{R})$, and the Riemannian product $\mathbb{S}^2\times\mathbb{R}$ and $\mathbb{H}^2\times\mathbb{R}$, where $\mathbb{S}^2$ and $\mathbb{H}^2$ are the 2-dimensional sphere and the hyperbolic plane, respectively. A crucial feature of these spaces is that they admit a Rimannian submersion onto a surface of constant Gaussian curvature, called the Hopf fibration.

In [1] A. Balmuş determined the parametric equations of all proper biharmonic curves on the Berger sphere $\mathbb{S}^3_\epsilon$ as curves in $\mathbb{R}^4$ and gave a geometric interpretation for those curves in the unit Euclidean sphere $\mathbb{S}^3$. In [6] the authors proved that any proper biharmonic curve in the Heisenberg group is an helix and gave their explicit parametrizations.
Also, in [4] are considered the proper biharmonic curves in the Bianchi-Cartan-Vranceanu spaces $\widetilde{\mathrm{SL}}(2,\mathbb{R})$, $SU(2)$, $\mathbb{S}^2\times\mathbb{R}$ and $\mathbb{H}^2\times\mathbb{R}$, proving that these curves are helices and giving their parametric equations.

In this paper we study the proper biharmonic curves in the special linear group $\mathrm{SL}(2,\mathbb{R})$ endowed with a suitable 1-parameter family g_τ of metrics that we shall describe in Section 2. Using the same technique given in [1] for the case of the Berger sphere, we conclude that the biharmonic curves of $\mathrm{SL}(2,\mathbb{R})$ makes a constant angle ϑ with the vector field tangent to the Hopf fibration and we prove the Theorem 3.4, which states that the differential equation

$$\gamma^{IV}+(b^2-2a)\,\gamma''+a^2\,\gamma=0$$

must be satisfied by any proper biharmonic curve in $\mathrm{SL}(2,\mathbb{R})$, as a curve in the pseudo-Euclidean $\mathbb{R}^4_2$, where a and b are real constants depending on ϑ and τ. We separate the study in three cases depending on the sign of the constant (b^2-4a) obtaining, in each case, the expressions of these curves as curves in $\mathbb{R}^4_2$.

2. Preliminaries

Let $\mathbb{R}^4_2$ denote the 4-dimensional pseudo-Euclidean space endowed with semi-definite inner product of signature $(2,2)$ given by

$$\langle v,w\rangle=v_1\,w_1+v_2\,w_2-v_3\,w_3-v_4\,w_4\,,\quad v,w\in\mathbb{R}^4.$$

We identify the special linear group with

$$\mathrm{SL}(2,\mathbb{R})=\{(z,w)\in\mathbb{C}^2\colon |z|^2-|w|^2=1\}=\{v\in\mathbb{R}^4_2\colon \langle v,v\rangle=1\}\subset\mathbb{R}^4_2$$

and we shall use the Lorentz model of the hyperbolic plane with constant Gauss curvature $-\tau$, $\tau>0$, that is

$$\mathbb{H}^2(-\tau)=\{(x,y,z)\in\mathbb{R}^3_1\colon x^2+y^2-z^2=-1/\tau\},$$

where $\mathbb{R}^3_1$ is the Minkowski 3-space. Then the Hopf map $\psi:\mathrm{SL}(2,\mathbb{R})\to\mathbb{H}^2(-\tau)$ given by

$$\psi(z,w)=\frac{1}{\sqrt{\tau}}\,(2z\bar{w},|z|^2+|w|^2)$$

is a submersion, with circular fibers, and if we put

$$X_1(z,w)=(iz,iw),\quad X_2(z,w)=(i\bar{w},i\bar{z}),\quad X_3(z,w)=(\bar{w},\bar{z}),$$

we have that X_1 is a vertical vector field while X_2, X_3 are horizontal. The vector X_1 is called the *Hopf vector field*.
We shall endow $\mathrm{SL}(2,\mathbb{R})$ with the 1-parameter family of metrics g_τ, $\tau > 0$, given by

$$g_\tau(X_i, X_j) = \delta_{ij}, \quad g_\tau(X_1, X_1) = \tau^2, \quad g_\tau(X_1, X_j) = 0, \quad i, j \in \{2, 3\},$$

which renders the Hopf map $\psi : (\mathrm{SL}(2,\mathbb{R}), g_\tau) \to \mathbb{H}^2(-\tau)$ a Riemannian submersion. With respect to the inner product in $\mathbb{R}^4_2$ the metric g_τ is given by

$$g_\tau(X, Y) = -\langle X, Y\rangle + (1+\tau^2)\langle X, X_1\rangle\langle Y, X_1\rangle\,. \tag{4}$$

From now on, we denote $(\mathrm{SL}(2,\mathbb{R}), g_\tau)$ with $\mathrm{SL}(2,\mathbb{R})_\tau$. Obviously

$$E_1 = -\tau^{-1} X_1, \quad E_2 = X_2, \quad E_3 = X_3, \tag{5}$$

is an orthonormal basis on $\mathrm{SL}(2,\mathbb{R})_\tau$. The Levi-Civita connection ∇^τ of $\mathrm{SL}(2,\mathbb{R})_\tau$ is given by:

$$\begin{aligned}
&\nabla^\tau{}_{E_1} E_1 = 0, \quad \nabla^\tau{}_{E_2} E_2 = 0, \quad \nabla^\tau{}_{E_3} E_3 = 0,\\
&\nabla^\tau{}_{E_1} E_2 = -\tau^{-1}(2+\tau^2) E_3, \quad \nabla^\tau{}_{E_1} E_3 = \tau^{-1}(2+\tau^2) E_2,\\
&\nabla^\tau{}_{E_2} E_1 = -\tau E_3, \quad \nabla^\tau{}_{E_3} E_1 = \tau E_2, \quad \nabla^\tau{}_{E_3} E_2 = -\tau E_1 = -\nabla^\tau{}_{E_2} E_3.
\end{aligned} \tag{6}$$

Using the conventions

$$\mathrm{R}(X, Y)Z = \nabla^\tau{}_X \nabla^\tau{}_Y Z - \nabla^\tau{}_Y \nabla^\tau{}_X Z - \nabla^\tau{}_{[X,Y]} Z$$

and

$$\mathrm{R}(X, Y, W, Z) = g_\tau(\mathrm{R}(X, Y)Z, W),$$

the nonzero components of the Riemannian curvature are

$$\mathrm{R}_{1212} = \tau^2, \qquad \mathrm{R}_{1313} = \tau^2, \qquad \mathrm{R}_{2323} = -(4 + 3\tau^2), \tag{7}$$

where $\mathrm{R}_{ijkl} = \mathrm{R}(E_i, E_j, E_k, E_l)$.

Finally, we recall that the isometry group of $\mathrm{SL}(2,\mathbb{R})_\tau$ is the 4-dimensional indefinite unitary group $\mathrm{U}_1(2)$ that can be identified with:

$$\mathrm{U}_1(2) = \{A \in \mathrm{O}_2(4) \colon AJ_1 = \pm J_1 A\}\,,$$

where J_1 is the complex structure of $\mathbb{R}^4$ defined by

$$J_1 = \begin{pmatrix} J & 0 \\ 0 & J \end{pmatrix}, \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

while

$$\mathrm{O}_2(4) = \{A \in \mathrm{GL}(4,\mathbb{R}) \colon A^t = \epsilon\, A^{-1}\, \epsilon\}, \qquad \epsilon = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

is the indefinite orthogonal group.

3. Biharmonic curves in $\mathrm{SL}(2,\mathbb{R})_\tau$

Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau$ be a differentiable curve parametrized by arc length and let $\{T, N, B\}$ be the orthonormal frame field tangent to $\mathrm{SL}(2,\mathbb{R})_\tau$ along $\gamma(s)$ defined as follows: we denote by T the unit vector field $\gamma'(s)$ tangent to $\gamma(s)$, by N the unit vector field in the direction of $\nabla^\tau{}_T T$ normal to γ, and we choose B so that $\{T, N, B\}$ is a positive oriented orthonormal basis. Then we have the following Frenet equations

$$\begin{cases} \nabla^\tau{}_T T = k_1 N, \\ \nabla^\tau{}_T N = -k_1 T + k_2 B, \\ \nabla^\tau{}_T B = -k_2 N, \end{cases} \tag{8}$$

where $k_1 = |\nabla^\tau{}_T T|$ is the geodesic curvature of γ and k_2 its geodesic torsion.

Theorem 3.1. *Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau$ be a curve parametrized by arc length. Then γ is proper biharmonic if and only if*

$$\begin{cases} k_1 = \text{constant} \neq 0, \\ k_1^2 + k_2^2 = \tau^2 - 4\,(1+\tau^2)\,B_1^2, \\ k_2' = -4(1+\tau^2)\,N_1\,B_1. \end{cases} \tag{9}$$

Proof. Consider a curve $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau$ parametrized by arc length. In this case the equation (2) becomes

$$(\nabla^\tau{}_T)^3 T - \mathrm{R}(T, \nabla^\tau{}_T T)T = 0. \tag{10}$$

Using the Frenet equations into (10), we obtain the conditions

$$\begin{cases} k_1 = \text{constant} \neq 0, \\ k_1^2 + k_2^2 = \mathrm{R}(T,N,T,N), \\ k_2' = -\,\mathrm{R}(T,N,T,B). \end{cases} \tag{11}$$

Writing

$$T = \sum_{i=1}^{3} T_i\,E_i, \qquad N = \sum_{i=1}^{3} N_i\,E_i, \qquad B = \sum_{i=1}^{3} B_i\,E_i, \tag{12}$$

and using (7), we have that

$$\begin{aligned} \mathrm{R}(T,N,T,N) &= \tau^2 - 4(1+\tau^2)B_1^2, \\ \mathrm{R}(T,N,T,B) &= 4(1+\tau^2)N_1\,B_1. \end{aligned}$$

□

Proposition 3.2. *If $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau$ is a proper biharmonic curve parametrized by arc length, then its geodesic curvature and torsion are constants.*

Proof. From the Frenet equations it results that

$$g_\tau(\nabla^\tau{}_T B, E_1) = -g_\tau(k_2 N, E_1) = -k_2\,N_1.$$

On the other hand, using (6), we get

$$\begin{aligned} g_\tau(\nabla^\tau{}_T B, E_1) &= g_\tau(B_1' E_1 + T_2 B_3 \nabla^\tau{}_{E_2} E_3 + T_3 B_2 \nabla^\tau{}_{E_3} E_2, E_1) \\ &= B_1' + \tau(T_2 B_3 - T_3 B_2) \\ &= B_1' - \tau N_1. \end{aligned}$$

Combining these two equations, we have

$$B_1' = (\tau - k_2) N_1. \tag{13}$$

Now, using (9) we obtain

$$k_2 k_2' = -4(1+\tau^2) B_1 B_1'. \tag{14}$$

From (13) and (14) it results that $(\tau - 2k_2)B_1 N_1 = 0$. Therefore, we have two possibilities: $B_1 N_1 = 0$ that, together with (9), implies $k_2' = 0$; or $k_2 = \frac{\tau}{2}$. So k_2 is constant. □

Proposition 3.3. *If $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau$ is a proper biharmonic curve parametrized by arc length, then it makes a constant angle with the Hopf vector field E_1 and its tangent vector field can be writen as*

$$\gamma'(s) = T = \cos\vartheta\, E_1 + \sin\vartheta \sin\beta(s)\, E_2 + \sin\vartheta \cos\beta(s)\, E_3, \tag{15}$$

where $\vartheta \in (0, \pi/2]$ and $\beta : I \to \mathbb{R}$ is a smooth function.

Proof. First we note that $B_1 \neq 0$. Indeed if $B_1 = 0$ and $N_1 = 0$, then the curve is the integral curve of the vector field E_1 and it is a geodesic. Moreover, if $B_1 = 0$ and $N_1 \neq 0$, from (13) we get $k_2 = \tau$ that, together with the second equation of (9), gives $k_1 = 0$.
Since $B_1 \neq 0$, the third equation of (9) and the Proposition 3.2 implies $N_1 = 0$. Now, using the equations (6) and (8) we obtain

$$k_1 N_1 = g_\tau(\nabla^\tau{}_T T, E_1) = T_1'.$$

We conclude that $T_1 =$ constant and we obtain the expression (15). □

Using the previous result we have the following

Theorem 3.4. *Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau \subset \mathbb{R}^4_2$ be a curve parametrized by arc length. Then γ is proper biharmonic if and only if, as a curve in $\mathbb{R}^4_2$, satisfies*

$$\gamma^{IV} + (b^2 - 2a)\gamma'' + a^2 \gamma = 0, \tag{16}$$

where a and b are the constants given by:

$$\begin{cases} a = \dfrac{1}{2}(-\tau^{-2} + 1 - (1+\tau^{-2})\cos 2\vartheta) - \tau^{-1}\cos\vartheta\, \beta', \\ b = \beta' = -\tau^{-1}(2+\tau^2)\cos\vartheta \pm \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)}, \end{cases} \tag{17}$$

with

$$\frac{4(1+\tau^2)}{(4+5\tau^2)} \leq \cos^2\vartheta < 1.$$

Proof. Writing

$$\gamma(s) = (x_1(s), x_2(s), x_3(s), x_4(s)),$$

from (15) we have that the coordinates functions of γ in $\mathbb{R}^4_2$ satisfies

$$\begin{cases} x_1' = \tau^{-1}\cos\vartheta\, x_2 + \sin\vartheta\cos\beta\, x_3 + \sin\vartheta\sin\beta\, x_4, \\ x_2' = -\tau^{-1}\cos\vartheta\, x_1 + \sin\vartheta\sin\beta\, x_3 - \sin\vartheta\cos\beta\, x_4, \\ x_3' = \sin\vartheta\cos\beta\, x_1 + \sin\vartheta\sin\beta\, x_2 + \tau^{-1}\cos\vartheta\, x_4, \\ x_4' = \sin\vartheta\sin\beta\, x_1 - \sin\vartheta\cos\beta\, x_2 - \tau^{-1}\cos\vartheta\, x_3. \end{cases} \tag{18}$$

Deriving (18), it results that

$$\begin{cases} x_1'' = a\, x_1 - b\, x_2', \\ x_2'' = a\, x_2 + b\, x_1', \\ x_3'' = a\, x_3 - b\, x_4', \\ x_4'' = a\, x_4 + b\, x_3', \end{cases} \tag{19}$$

where

$$\begin{cases} a = \dfrac{1}{2}(-\tau^{-2} + 1 - (1+\tau^{-2})\cos 2\vartheta) - \tau^{-1}\cos\vartheta\,\beta', \\ b = \beta'. \end{cases}$$

Now, we shall prove that b is constant and we determine its expression. Computing $\nabla^\tau{}_T T$, using (15) and (6), the geodesic curvature and the normal vector field are given by

$$k_1 = \pm\sin\vartheta(\beta' + 2\tau^{-1}(1+\tau^2)\cos\vartheta), \qquad N = \pm(\cos\beta\, E_2 - \sin\beta\, E_3). \tag{20}$$

Then

$$\begin{aligned} B &= T \wedge N = \pm(-\sin\vartheta\, E_1 + \cos\vartheta\sin\beta\, E_2 + \cos\vartheta\cos\beta\, E_3), \\ k_2 &= g_\tau(\nabla^\tau{}_T N, B) = (\tau - \cos\vartheta(\beta' + 2\tau^{-1}(1+\tau^2)\cos\vartheta)). \end{aligned} \tag{21}$$

Substituting the expressions of k_1, k_2 and B_1 in the second equation of (9), it results that

$$\beta' = -\tau^{-1}(2+\tau^2)\cos\vartheta \pm \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)}.$$

Now deriving twice (19), and use (18), we obtain the equation (16). Also, as the curve γ is not harmonic, from (20), $\cos\vartheta \neq 1$.

□

Remark 3.5. Using (18) and (19), we find that:

$$\begin{array}{lll} \langle\gamma,\gamma\rangle = 1\,, & \langle\gamma',\gamma'\rangle = \tilde{B}, & \langle\gamma,\gamma'\rangle = 0, \\ \langle\gamma',\gamma''\rangle = 0\,, & \langle\gamma'',\gamma''\rangle = D\,, & \langle\gamma,\gamma''\rangle = -\tilde{B}, \\ \langle\gamma',\gamma'''\rangle = -D\,, & \langle\gamma'',\gamma'''\rangle = 0\,, & \langle\gamma,\gamma'''\rangle = 0, \\ \langle\gamma''',\gamma'''\rangle = E, & & \end{array} \tag{22}$$

where

$$\tilde{B} = (1+\tau^{-2})\cos^2\vartheta - 1, \qquad D = a^2 + b^2\tilde{B} + 2\,a\,b\,\tau^{-1}\cos\vartheta,$$
$$E = a\big(a - 2b^2\big)\tilde{B} + b^2 D - 2a^2 b\,\tau^{-1}\cos\vartheta.$$

In addition, as

$$J_1\gamma = X_{1|\gamma} = -\tau\, E_{1|\gamma},$$

using (15) and (19), we obtain the following identities

$$
\begin{aligned}
&\langle J_1\gamma, \gamma'\rangle = -\tau^{-1}\cos\vartheta,\\
&\langle J_1\gamma, \gamma''\rangle = 0\,,\\
&\langle J_1\gamma'', \gamma'\rangle = -a\,\tau^{-1}\,\cos\vartheta - b\,\tilde{B} := I,\\
&\langle J_1\gamma', \gamma'''\rangle = 0\,,\\
&\langle J_1\gamma', \gamma''\rangle + \langle J_1\gamma, \gamma'''\rangle = 0\,,\\
&\langle J_1\gamma'', \gamma'''\rangle + \langle J_1\gamma', \gamma^{IV}\rangle = 0\,.
\end{aligned}
\tag{23}
$$

To determine the expression of the position vector of γ in $\mathbb{R}^4_2$, we integrate (16), dividing the study in three cases, according to the three possibilities:

(i) $b^2 = 4a$;
(ii) $b^2 > 4a$;
(iii) $b^2 < 4a$.

4. The case $b^2 = 4a$

Theorem 4.1. *Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau \subset \mathbb{R}^4_2$ be a proper biharmonic curve parametrized by arc length such that $b^2 = 4a$. Then*

$$
b = -\tau^{-1}(2+\tau^2)\cos\vartheta + \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)},
\tag{24}
$$

with

$$
\cos^2\vartheta = \frac{(2+\tau^2)^2}{4+5\tau^2+\tau^4}.
$$

Also,

$$
\begin{aligned}
\gamma(s) =& A\Big(\cos(\sqrt{a}\,s) + g_{14}\,s\,\sin(\sqrt{a}\,s), -\sin(\sqrt{a}\,s) + g_{14}\,s\,\cos(\sqrt{a}\,s),\\
&- g_{14}\,s\,\cos(\sqrt{a}\,s),\, g_{14}\,s\,\sin(\sqrt{a}\,s)\Big),
\end{aligned}
\tag{25}
$$

where g_{14} is the constant, given by

$$
g_{14} = \frac{\tau}{\sqrt{4+5\tau^2+\tau^4}}
$$

and $A \in \mathrm{O}_2(4)$ is a 4×4 indefinite orthogonal matrix which commutes with J_1.

Proof. As $b^2 = 4a$, the differential equation (16) turns

$$
\gamma^{IV}(s) + 2a\,\gamma''(s) + a^2\gamma(s) = 0.
\tag{26}
$$

Integrating (26) we have

$$
\gamma(s) = \cos(\sqrt{a}\,s)\,g_1 + \sin(\sqrt{a}\,s)\,g_2 + s\,\cos(\sqrt{a}\,s)\,g_3 + s\,\sin(\sqrt{a}\,s)\,g_4,
\tag{27}
$$

where g_1, g_2, g_3 and g_4 are constant vectors of $\mathbb{R}^4_2$.

A direct calculation shows that $b^2 = 4a$ occurs in two cases: for $\vartheta = 0$ and for

$$
\cos^2\vartheta = \frac{(2+\tau^2)^2}{4+5\tau^2+\tau^4},
$$

and in both cases b must have the expression given in (24). Since the first case produces harmonic curves, we study only the second one.

Using the relations (22), we get

$$\begin{aligned}
&\langle g_1, g_1\rangle = \langle g_2, g_2\rangle = 1,\\
&\langle g_3, g_3\rangle = \langle g_4, g_4\rangle = 0,\\
&\langle g_1, g_4\rangle = -\langle g_2, g_3\rangle = \frac{\tau}{\sqrt{4+5\tau^2+\tau^4}},\\
&\langle g_1, g_2\rangle = \langle g_1, g_3\rangle = \langle g_2, g_4\rangle = \langle g_3, g_4\rangle = 0,
\end{aligned} \tag{28}$$

whereas (23) yields

$$\begin{aligned}
&\langle J_1 g_1, g_2\rangle = -1,\\
&\langle J_1 g_2, g_4\rangle = \langle J_1 g_1, g_3\rangle = \frac{\tau}{\sqrt{4+5\tau^2+\tau^4}},\\
&\langle J_1 g_1, g_4\rangle = \langle J_1 g_2, g_3\rangle = \langle J_1 g_3, g_4\rangle = 0.
\end{aligned} \tag{29}$$

Now, putting

$$\begin{cases}
e_1 = g_1,\\
e_2 = g_2,\\
e_3 = \dfrac{g_3}{\langle g_2, g_3\rangle} - g_2,\\
e_4 = \dfrac{g_4}{\langle g_1, g_4\rangle} - g_1,
\end{cases}$$

we have that $\{e_i\}$ is an orthonormal basis of $\mathbb{R}^4_2$ that satisfies:

$$\begin{aligned}
&\langle J_1 e_1, e_2\rangle = \langle J_1 e_3, e_4\rangle = -1,\\
&\langle J_1 e_1, e_3\rangle = \langle J_1 e_1, e_4\rangle = \langle J_1 e_2, e_3\rangle = \langle J_1 e_2, e_4\rangle = 0.
\end{aligned}$$

We conclude that $e_2 = -J_1 e_1$ and $e_4 = J_1 e_3$. So if we consider the orthonormal basis $\{\tilde{E}_i\}_{i=1}^4$ of $\mathbb{R}^4_2$ given by

$$\tilde{E}_1 = (1,0,0,0)\,, \quad \tilde{E}_2 = (0,-1,0,0)\,, \quad \tilde{E}_3 = (0,0,1,0)\,, \quad \tilde{E}_4 = (0,0,0,1)\,,$$

there must exists a matrix $A \in \mathrm{O}_2(4)$, with $J_1\, A = A\, J_1$ such that $e_i = A\,\tilde{E}_i$, $i \in \{1,2,3,4\}$. Finally, putting $\langle g_1, g_4\rangle = g_{14}$, we can rewrite (27) as (25). □

5. The case $b^2 > 4a$

Theorem 5.1. *Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau \subset \mathbb{R}^4_2$ be a proper biharmonic curve parametrized by arc length, such that $b^2 > 4a$. Then there are two possibilities:*

(i)

$$b = -\tau^{-1}(2+\tau^2)\cos\vartheta + \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)}$$

and

$$\frac{4(1+\tau^2)}{(4+5\tau^2)} \le \cos^2\vartheta < \frac{(2+\tau^2)^2}{4+5\tau^2+\tau^4};$$

(ii)

$$b = -\tau^{-1}(2+\tau^2)\cos\vartheta - \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)}$$

and

$$\frac{4(1+\tau^2)}{(4+5\tau^2)} \leq \cos^2\vartheta.$$

In both cases, the expression of γ *as a curve in* $\mathbb{R}^4_2$ *is*

$$\gamma(s) = A\big(\sqrt{C_{33}}\cos(\alpha_2\, s)\,,\ \sqrt{C_{33}}\sin(\alpha_2\, s)\,,\ \sqrt{-C_{11}}\cos(\alpha_1\, s)\,,\ \sqrt{-C_{11}}\sin(\alpha_1\, s)\big)\,, \tag{30}$$

where

$$\alpha_{1,2} = \sqrt{\frac{(b^2-2a) \pm \sqrt{b^2(b^2-4a)}}{2}}$$

and

$$C_{11} = \frac{\tilde{B} - \alpha_2^2}{\alpha_1^2 - \alpha_2^2}, \qquad C_{33} = \frac{-\tilde{B} + \alpha_1^2}{\alpha_1^2 - \alpha_2^2},$$

are real constants and $A \in \mathrm{O}_2(4)$ *is a* 4×4 *indefinite orthogonal matrix anticommuting with* J_1.

Proof. First, observe that the condition $b^2 > 4a$ gives the two possibilities (i) and (ii). Also a direct integration of (16), gives the solution

$$\gamma(s) = \cos(\alpha_1\, s)\, C_1 + \sin(\alpha_1\, s)\, C_2 + \cos(\alpha_2\, s)\, C_3 + \sin(\alpha_2\, s)\, C_4,$$

where

$$\alpha_{1,2} = \sqrt{\frac{(b^2-2a) \pm \sqrt{b^2(b^2-4a)}}{2}}$$

are real constants, while the C_i, $i \in \{1,2,3,4\}$, are constants vectors in $\mathbb{R}^4_2$.

Putting $C_{ij} = \langle C_i, C_j\rangle$, and evaluating the relations (22) in $s=0$, we obtain:

$$C_{11} + C_{33} + 2C_{13} = 1, \tag{31}$$

$$\alpha_1^2\, C_{22} + \alpha_2^2\, C_{44} + 2\alpha_1\alpha_2\, C_{24} = \tilde{B}, \tag{32}$$

$$\alpha_1\, C_{12} + \alpha_2\, C_{14} + \alpha_1\, C_{23} + \alpha_2\, C_{34} = 0, \tag{33}$$

$$\alpha_1^3\, C_{12} + \alpha_1\alpha_2^2\, C_{23} + \alpha_1^2\alpha_2\, C_{14} + \alpha_2^3 C_{34} = 0, \tag{34}$$

$$\alpha_1^4\, C_{11} + \alpha_2^4\, C_{33} + 2\alpha_1^2\alpha_2^2\, C_{13} = D, \tag{35}$$

$$\alpha_1^2\, C_{11} + \alpha_2^2\, C_{33} + (\alpha_1^2 + \alpha_2^2)\, C_{13} = \tilde{B}, \tag{36}$$

$$\alpha_1^4\, C_{22} + (\alpha_1^3\alpha_2\ + \alpha_1\alpha_2^3)\, C_{24} + \alpha_2^4\, C_{44} = D, \tag{37}$$

$$\alpha_1^5\, C_{12} + \alpha_1^3\alpha_2^2\, C_{23} + \alpha_1^2\alpha_2^3\, C_{14} + \alpha_2^5\, C_{34} = 0, \tag{38}$$

$$\alpha_1^3\, C_{12} + \alpha_1^3\, C_{23} + \alpha_2^3\, C_{14} + \alpha_2^3\, C_{34} = 0, \tag{39}$$

$$\alpha_1^6\, C_{22} + \alpha_2^6\, C_{44} + 2\alpha_1^3\alpha_2^3\, C_{24} = E. \tag{40}$$

From (33), (34), (38), (39), it follows that

$$C_{12} = C_{14} = C_{23} = C_{34} = 0.$$

Also, from (31), (35) and (36), we obtain

$$C_{11} = \frac{\tilde{B} - \alpha_2^2}{\alpha_1^2 - \alpha_2^2}, \qquad C_{13} = 0, \qquad C_{33} = \frac{-\tilde{B} + \alpha_1^2}{\alpha_1^2 - \alpha_2^2}.$$

Finally, using (32), (37) and (40), we obtain

$$C_{22} = \frac{D - \tilde{B}\alpha_2^2}{\alpha_1^2(\alpha_1^2 - \alpha_2^2)}, \quad C_{24} = 0, \qquad C_{44} = \frac{-D + \tilde{B}\alpha_1^2}{\alpha_2^2(\alpha_1^2 - \alpha_2^2)}.$$

We observe that as

$$\frac{4(1+\tau^2)}{(4+5\tau^2)} \leq \cos^2\vartheta,$$

then

$$C_{11} = C_{22} < 0, \qquad C_{33} = C_{44} > 0.$$

Since $\{C_i\}_{i=1}^4$ are mutually orthogonal and

$$||C_1|| = ||C_2|| = \sqrt{-C_{11}}, \qquad ||C_3|| = ||C_4|| = \sqrt{C_{33}},$$

we obtain a pseudo-orthonormal basis of $\mathbb{R}_2^4$ putting $e_i = C_i/||C_i||$, $i \in \{1,2,3,4\}$, and we can write:

(41) $\gamma(s) = \sqrt{-C_{11}}\,\big(\cos(\alpha_1 s)e_1 + \sin(\alpha_1 s)e_2\big) + \sqrt{C_{33}}\,\big(\cos(\alpha_2 s)e_3 + \sin(\alpha_2 s)e_4\big)$.

Now, evaluating in $s = 0$ the identities (23), we have:

$$\begin{aligned} &\alpha_2\, C_{33}\langle J_1e_3, e_4\rangle - \alpha_1 C_{11}\langle J_1e_1, e_2\rangle \\ &+ \sqrt{-C_{11}C_{33}}\,(\alpha_1\langle J_1e_3, e_2\rangle + \alpha_2\langle J_1e_1, e_4\rangle) = -\tau^{-1}\cos\vartheta, \\ &\langle J_1e_1, e_3\rangle = 0\,, \end{aligned} \tag{42}$$

$$\begin{aligned} &\alpha_2^3\, C_{33}\langle J_1e_3, e_4\rangle - \alpha_1^3\, C_{11}\langle J_1e_1, e_2\rangle \\ &+ \sqrt{-C_{11}C_{33}}\,(\alpha_1\alpha_2^2\langle J_1e_3, e_2\rangle + \alpha_1^2\alpha_2\langle J_1e_1, e_4\rangle) = -I, \\ &\langle J_1e_2, e_4\rangle = 0\,, \end{aligned} \tag{43}$$

$$\alpha_1\langle J_1e_2, e_3\rangle + \alpha_2\langle J_1e_1, e_4\rangle = 0\,, \tag{44}$$

$$\alpha_2\langle J_1e_2, e_3\rangle + \alpha_1\langle J_1e_1, e_4\rangle = 0\,. \tag{45}$$

We point out that to obtain the previous identities we have divided by $\alpha_1^2 - \alpha_2^2 = \sqrt{b^2(b^2-4a)}$ which is different from zero. From (44) and (45), taking into account the $\alpha_1^2 - \alpha_2^2 \neq 0$, it results that

$$\langle J_1e_3, e_2\rangle = 0\,, \qquad \langle J_1e_1, e_4\rangle = 0\,. \tag{46}$$

Then, $J_1e_1 = \pm e_2$ and $J_1e_3 = \pm e_4$. So, the position vector of γ is given by

(47) $\gamma(s) = \sqrt{-C_{11}}\,\big(\cos(\alpha_1 s)e_1 \pm \sin(\alpha_1 s)J_1e_1\big) + \sqrt{C_{33}}\,\big(\cos(\alpha_2 s)e_3 \pm \sin(\alpha_2 s)J_1e_3\big)$.

If we use (19) for $s = 0$, we get $J_1e_1 = -e_2$ and $J_1e_3 = -e_4$.

Then, if we fix the orthonormal basis of $\mathbb{R}^4_2$ given by

$$\bar{E}_1 = (0,0,1,0)\,, \quad \bar{E}_2 = (0,0,0,1)\,, \quad \bar{E}_3 = (1,0,0,0)\,, \quad \bar{E}_4 = (0,1,0,0)\,,$$

there must exists a matrix $A \in \mathrm{O}_2(4)$, with $J_1\,A = -A\,J_1$, such that $e_i = A\,\bar{E}_i$. □

6. The case $b^2 < 4a$

Theorem 6.1. *Let $\gamma : I \to \mathrm{SL}(2,\mathbb{R})_\tau \subset \mathbb{R}^4_2$ be a proper biharmonic curve parametrized by arc length, such that $b^2 < 4a$. Then*

$$b = -\tau^{-1}(2+\tau^2)\cos\vartheta + \sqrt{(4+5\tau^2)\cos^2\vartheta - 4(1+\tau^2)}, \tag{48}$$

$$\frac{(2+\tau^2)^2}{4+5\tau^2+\tau^4} < \cos^2\vartheta < 1, \tag{49}$$

and the expression of γ as a curve in $\mathbb{R}^4_2$ is

$$\begin{aligned}\gamma(s) =& A\Big(\cos\big(\frac{b}{2}\,s\big)\cosh(\mu\,s) + w_{14}\sin\big(\frac{b}{2}\,s\big)\sinh(\mu\,s)\,,\\ &\sin\big(\frac{b}{2}\,s\big)\cosh(\mu\,s) - w_{14}\cos\big(\frac{b}{2}\,s\big)\sinh(\mu\,s)\,,\\ &\cos\big(\frac{b}{2}\,s\big)\sinh(\mu\,s)\sqrt{1+w_{14}^2}\,,\ \sin\big(\frac{b}{2}\,s\big)\sinh(\mu\,s)\sqrt{1+w_{14}^2}\Big),\end{aligned} \tag{50}$$

where

$$\mu = \frac{\sqrt{4a-b^2}}{2}, \qquad w_{14} = \frac{b\tau + 2\cos\vartheta}{2\tau\mu}$$

are real constants and $A \in \mathrm{O}_2(4)$ is a 4×4 indefinite orthogonal matrix commuting with J_1.

Proof. From $b^2 < 4a$, it results that b ig given by (48) and θ satisfies (49). Also a direct integration of (16), gives

$$\gamma(s) = \cos\big(\frac{b}{2}\,s\big)\,\big(\cosh(\mu\,s)\,w_1 + \sinh(\mu\,s)\,w_3\big) + \sin\big(\frac{b}{2}\,s\big)\,\big(\cosh(\mu\,s)\,w_2 + \sinh(\mu\,s)\,w_4\big),$$

where

$$\mu = \frac{\sqrt{4a-b^2}}{2},$$

while the w_i, $i \in \{1,2,3,4\}$, are constant vectors in $\mathbb{R}^4_2$. If $w_{ij} := \langle w_i, w_j\rangle$, evaluating the relations (22) in $s = 0$, we obtain

$$w_{11} = 1, \tag{51}$$

$$\frac{b^2}{4}\,w_{22} + \mu^2\,w_{33} + \mu\,b\,w_{23} = \tilde{B}, \tag{52}$$

$$\frac{b}{2}\,w_{12} + \mu\,w_{13} = 0, \tag{53}$$

$$\frac{b}{2}\left(\mu^2 - \frac{b^2}{4}\right)w_{12} + \mu^2\,b\,w_{34} + \mu\,\frac{b^2}{2}\,w_{24} + \mu\left(\mu^2 - \frac{b^2}{4}\right)w_{13} = 0, \tag{54}$$

$$\left(\mu^2 - \frac{b^2}{4}\right)^2 w_{11} + \mu^2\,b^2\,w_{44} + 2\mu\,b\left(\mu^2 - \frac{b^2}{4}\right)w_{14} = D, \tag{55}$$

$$\left(\mu^2 - \frac{b^2}{4}\right) w_{11} + \mu\, b\, w_{14} = -\tilde{B}, \tag{56}$$

$$\frac{b^2}{4}\left(3\mu^2 - \frac{b^2}{4}\right) w_{22} + \mu^2\left(\mu^2 - 3\frac{b^2}{4}\right) w_{33} + \mu\,\frac{b}{2}\,(4\mu^2 - b^2)\, w_{23} = -D, \tag{57}$$

$$\begin{aligned} &\frac{b}{2}\left(3\mu^2 - \frac{b^2}{4}\right)\left(\mu^2 - \frac{b^2}{4}\right) w_{12} + b\,\mu^2\left(\mu^2 - 3\frac{b^2}{4}\right) w_{34} \\ &+ \mu\left(\mu^2 - 3\frac{b^2}{4}\right)\left(\mu^2 - \frac{b^2}{4}\right) w_{13} + \mu\,\frac{b^2}{2}\left(3\mu^2 - \frac{b^2}{4}\right) w_{24} = 0, \end{aligned} \tag{58}$$

$$\frac{b}{2}\left(3\mu^2 - \frac{b^2}{4}\right) w_{12} + \mu\left(\mu^2 - 3\frac{b^2}{4}\right) w_{13} = 0, \tag{59}$$

$$\begin{aligned} &\frac{b^2}{4}\left(3\mu^2 - \frac{b^2}{4}\right)^2 w_{22} + \mu^2\left(\mu^2 - 3\frac{b^2}{4}\right)^2 w_{33} \\ &+ \mu\, b\left(3\mu^2 - \frac{b^2}{4}\right)\left(\mu^2 - 3\frac{b^2}{4}\right) w_{23} = E. \end{aligned} \tag{60}$$

From (51), (55) and (56), it follows that

$$w_{11} = -w_{44} = 1, \qquad w_{14} = \frac{b\tau + 2\cos\vartheta}{2\tau\mu}.$$

Also, from (53) and (59), we obtain

$$w_{12} = w_{13} = 0$$

and, therefore, from (54) and (58),

$$w_{24} = w_{34} = 0.$$

Moreover, using (52), (57) and (60), we get

$$w_{22} = -w_{33} = 1, \qquad w_{23} = -\frac{b\tau + 2\cos\vartheta}{2\tau\mu}.$$

Then we can define the following pseudo-orthonormal basis in $\mathbb{R}^4_2$:

$$\begin{cases} e_1 = w_1, \\ e_2 = w_2, \\ e_3 = \dfrac{w_3 + w_{14}\, w_2}{\sqrt{1 + w_{14}^2}}, \\ e_4 = \dfrac{w_4 - w_{14}\, w_1}{\sqrt{1 + w_{14}^2}}, \end{cases}$$

with $\langle e_1, e_1\rangle = 1 = \langle e_2, e_2\rangle$ and $\langle e_3, e_3\rangle = -1 = \langle e_4, e_4\rangle$.

Evaluating the identities (23) in $s=0$, and taking into account that:

$$\begin{aligned}
&\gamma(0) = w_1\,,\\
&\gamma'(0) = \frac{b}{2}\,w_2 + \mu\,w_3\,,\\
&\gamma''(0) = \Big(\mu^2 - \frac{b^2}{4}\Big)\,w_1 + \mu\,b\,w_4\,,\\
&\gamma'''(0) = \frac{b}{2}\,\Big(3\mu^2 - \frac{b^2}{4}\Big)\,w_2 + \mu\,\Big(\mu^2 - \frac{3}{4}b^2\Big)\,w_3\,,\\
&\gamma^{IV}(0) = \Big(\mu^4 - \frac{3}{2}\mu^2\,b^2 + \frac{b^4}{16}\Big)\,w_1 + 2\mu\,b\,\Big(\mu^2 - \frac{b^2}{4}\Big)\,w_4\,,
\end{aligned}$$

we conclude that

$$\begin{aligned}
&\langle J_1 w_1, w_2\rangle = -\langle J_1 w_3, w_4\rangle = 1,\\
&\langle J_1 w_3, w_2\rangle = \langle J_1 w_1, w_4\rangle = 0,\\
&\langle J_1 w_1, w_3\rangle = \langle J_1 w_2, w_4\rangle = -w_{14}.
\end{aligned}$$

Then,

$$\begin{aligned}
&\langle J_1 e_1, e_2\rangle = -\langle J_1 e_3, e_4\rangle = 1,\\
&\langle J_1 e_1, e_4\rangle = \langle J_1 e_1, e_3\rangle = \langle J_1 e_2, e_3\rangle = \langle J_1 e_2, e_4\rangle = 0.
\end{aligned}$$

Therefore, we obtain that

$$J_1 e_1 = e_2, \qquad J_1 e_3 = e_4.$$

Consequently, if we consider the orthonormal basis $\{E_i\}_{i=1}^4$ of $\mathbb{R}^4_2$ given by

$$E_1 = (1,0,0,0)\,, \quad E_2 = (0,1,0,0)\,, \quad E_3 = (0,0,1,0)\,, \quad E_4 = (0,0,0,1)\,,$$

there must exists $A \in \mathrm{O}_2(4)$, with $J_1 A = A J_1$, such that $e_i = A\,E_i$, $i \in \{1,2,3,4\}$. □

References

[1] A. Balmuş. On the biharmonic curves of the Euclidean and Berger 3-dimensional spheres. *Sci. Ann. Univ. Agric. Sci. Vet. Med.* 47 (2004), 87–96.

[2] R. Caddeo, S. Montaldo, C. Oniciuc. Biharmonic submanifolds of $\mathbb{S}^3$. *Internat. J. Math.* 12 (2001), 867–876.

[3] R. Caddeo, S. Montaldo, C. Oniciuc. Biharmonic submanifolds in spheres. *Israel J. Math.* 130 (2002), 109–123.

[4] R. Caddeo, S. Montaldo, C. Oniciuc, P. Piu. The Euler-Lagrange method for biharmonic curves. *Mediterr. J. Math.* 3 (2006), no. 3-4, 449–465.

[5] R. Caddeo, S. Montaldo, P. Piu. Biharmonic curves on a surface. *Rendiconti di Matematica e delle sue Applicazioni* Serie VII, Volume 21, Roma (2001), 143–157.

[6] R. Caddeo, P. Piu, C. Oniciuc. Explicit formulas for non-geodesic biharmonic curves of the Heisenberg group. *Ren. Mat. Univ. Politc. Torino* 62 (2004), no. 3, 265–277.

[7] I. Dimitric. Submanifolds of $\mathbb{E}^m$ with harmonic mean curvature vetor. *Bull. Inst. Math. Acad. Sinica* 20 (1992), 53–65.

[8] J. Eells and L. Lemaire. Selected topics in harmonic maps. *Conf. Board Math. Sci.* 50 (1983).

[9] G.Y Jiang. 2-harmonic isometric immersions between Riemannian manifolds. *Chinese Ann. Math. Ser* A 7 (1986), no. 2, 130–144.

[10] G.Y Jiang. 2-harmonic maps and their first and second variational formulas. *Chinese Ann. Math. Ser* A 7 (1986), no. 4, 389–402.

Departamento de Matemática, C.P. 668, ICMC, USP, 13560-970, São Carlos, SP, Brasil
E-mail address: onnis@icmc.usp.br

Departamento de Matemática, C.P. 668, ICMC, USP, 13560-970, São Carlos, SP, Brasil
E-mail address: apoenapp@icmc.usp.br